



Applying a new ensemble approach to estimating stock status of marine fisheries around the world

Estimating global fisheries status

Rosenberg, Andrew A.; Kleisner, Kristin M.; Afflerbach, Jamie; Anderson, Sean C.; Dickey-Collas, Mark; Cooper, Andrew B.; Fogarty, Michael J.; Fulton, Elizabeth A.; Gutiérrez, Nicolás L.; Hyde, Kimberly J.W.

Total number of authors:
23

Published in:
Conservation Letters

Link to article, DOI:
[10.1111/conl.12363](https://doi.org/10.1111/conl.12363)

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Rosenberg, A. A., Kleisner, K. M., Afflerbach, J., Anderson, S. C., Dickey-Collas, M., Cooper, A. B., Fogarty, M. J., Fulton, E. A., Gutiérrez, N. L., Hyde, K. J. W., Jardim, E., Jensen, O. P., Kristiansen, T., Longo, C., Mente-Vera, C. V., Minto, C., Mosqueira, I., Osio, G. C., Ovando, D., ... Ye, Y. (2018). Applying a new ensemble approach to estimating stock status of marine fisheries around the world: Estimating global fisheries status. *Conservation Letters*, 11(1), [e12363]. <https://doi.org/10.1111/conl.12363>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LETTER

Applying a New Ensemble Approach to Estimating Stock Status of Marine Fisheries Around the World

Andrew A. Rosenberg¹, Kristin M. Kleisner², Jamie Afflerbach³, Sean C. Anderson⁴, Mark Dickey-Collas^{5,6}, Andrew B. Cooper⁷, Michael J. Fogarty⁸, Elizabeth A. Fulton^{9,10}, Nicolás L. Gutiérrez¹¹, Kimberly J. W. Hyde¹², Ernesto Jardim¹³, Olaf P. Jensen¹⁴, Trond Kristiansen¹⁵, Catherine Longo¹⁶, Carolina V. Minto-Vera¹⁷, Cólín Minto¹⁸, Iago Mosqueira¹³, Giacomo Chato Osio¹³, Daniel Ovando¹⁹, Elizabeth R. Selig^{20,21}, James T. Thorson²², Jessica C. Walsh⁷, & Yimin Ye¹¹

¹ Center for Science and Democracy, Union of Concerned Scientists, 2 Brattle Square, Cambridge, MA 02138, USA

² Environmental Defense Fund, 18 Tremont St., Ste. 850, Boston, MA 02108, USA

³ National Center for Ecological Analysis and Synthesis, University of California Santa Barbara, 735 State Street, Santa Barbara, CA 93101, USA

⁴ School of Aquatic and Fishery Sciences, University of Washington, Box 455020, Seattle, WA 98195, USA

⁵ International Council for the Exploration of the Seas, H. C. Andersens Boulevard 44–46, DK 1553, Copenhagen, Denmark

⁶ National Institute of Aquatic Resources, Danish Technical University, Jægersborg Alle 1, 2920 Charlottenlund, Denmark

⁷ School of Resource and Environmental Management, Simon Fraser University, Burnaby, BC, V5A 1S6 Canada

⁸ Northeast Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration 166 Water St., Woods Hole, MA 02543, USA

⁹ CSIRO Oceans & Atmosphere, Castray Esplanade, Hobart, Tasmania 7001, Australia

¹⁰ Centre Marine Socioecology, University of Tasmania, Hobart, Tasmania 7001, Australia

¹¹ Food and Agriculture Organization, Via del Terme di Caracalla 00153, Rome, Italy

¹² Northeast Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Narragansett, RI 02882, USA

¹³ European Commission, DG Joint Research Center, Directorate D – Sustainable Resources, Unit D.02 Water and Marine Resources, Via Enrico Fermi 2749 21027 Ispra (VA), Italy

¹⁴ Department of Marine & Coastal Sciences, Rutgers University, 71 Dudley Rd., New Brunswick, NJ 08540, USA

¹⁵ Institute of Marine Research, P.O. Box 1870 Nordnes, Bergen 5817, Norway

¹⁶ Marine Stewardship Council, Marine House, 1 Snow Hill, London, EC1A 2DH UK

¹⁷ Nupelia-State University of Maringá, Oceanographic Institute, University of São Paulo São Paulo, Brazil

¹⁸ Galway-Mayo Institute of Technology, Marine and Freshwater Research Centre, Galway, Dublin Road, Ireland

¹⁹ Bren School of Environmental Science and Management, University of California, Santa Barbara, CA 93106, USA

²⁰ Conservation International, 2011 Crystal Drive, Suite 500, Arlington, VA 22202, USA

²¹ Norwegian Institute for Water Research, Gaustadalléen 21, 0349 Oslo, Norway

²² Fisheries Resource and Monitoring Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, 2725 Montlake Blvd. East, Seattle, WA 98112, USA

Keywords

Fisheries exploitation status; global fisheries; food security; ecosystem sustainability.

Correspondence

Andrew A. Rosenberg, Union of Concerned Scientists, 2 Brattle Square, Cambridge, MA 02138, USA. Tel: 617-301-8010.
E-mail: arosenberg@ucsusa.org

Received

11 July 2016

Accepted

12 March 2017

doi: 10.1111/conl.12363

Abstract

The exploitation status of marine fisheries stocks worldwide is of critical importance for food security, ecosystem conservation, and fishery sustainability. Applying a suite of data-limited methods to global catch data, combined through an ensemble modeling approach, we provide quantitative estimates of exploitation status for 785 fish stocks. Fifty-six percent (439 stocks) are below B_{MSY} and of these, 261 are estimated to be below 80% of the B_{MSY} level. While the 178 stocks above 80% of B_{MSY} are conventionally considered “fully exploited,” stocks staying at this level for many years, forego substantial yield. Our results enable managers to consider more detailed information than simply a categorization of stocks as “fully” or “over” exploited. Our approach is reproducible, allows consistent application to a broad range of stocks, and can be easily updated as new data become available. Applied on an ongoing basis, this approach can provide critical, more detailed information for resource management for more exploited fish stocks than currently available.

Introduction

The overexploitation of wild-capture marine fisheries directly impacts ecosystem health and the food security, livelihoods, and cultural identities of coastal communities worldwide (Garcia & Rosenberg 2010; Costello *et al.* 2012; FAO 2014; Inniss *et al.* 2016). The failure to sustainably manage some fisheries has resulted from many factors including deficient institutional capacity and inadequate data collection, leading to the lack of regular evaluations of stock exploitation status. Quantitative stock assessments are generally only available for relatively “data-rich,” commercially important stocks predominantly fished by developed countries. Recent studies of marine fisheries exploitation status in developed regions of the world have demonstrated that significant progress has been made in managing marine fisheries (e.g., Worm *et al.* 2009; Ricard *et al.* 2012; Fernandes & Cook 2013; Hilborn & Ovando 2014). However, much less is known about many smaller or lower valued stocks and stocks fished in both developed and developing regions of the world.

Recent estimates of the current state and future trajectory of marine stocks (Pauly *et al.* 1998; Worm *et al.* 2009; Garcia & Rosenberg 2010; Anderson *et al.* 2012; Costello *et al.* 2012, 1992; Thorson *et al.* 2012; FAO 2014) show that many of the world’s fisheries are below biologically sustainable biomass levels relative to widely accepted reference points. Stocks in data-limited regions are often in even poorer condition than stocks in data-rich regions (e.g., Costello *et al.* 2012). For these important marine resources, managers may at best only have a gross characterization of status, which provides little guidance for new policy action. Here, we apply a consistent, easily repeatable method for estimating more detailed quantitative estimates of stock status across both data-rich and data-limited fisheries.

Currently, there are two primary sources of information used to estimate the status of fish stocks (Thorson *et al.* 2012). The first source comes from abundance estimates derived from analytical stock assessments, developed by fisheries scientists around the world over the last 60 years (Beverton & Holt 1957; Hilborn & Walters 1992; Quinn & Deriso 1999). The data requirements for these sorts of quantitative stock assessment are substantial and are usually restricted to commercial fisheries where there is the analytical capacity to apply complex stock assessment models. There is a comprehensive compilation of fish stock assessments maintained in the RAM Legacy database (<http://ramlegacy.org>, Ricard *et al.* 2012). Most stocks included in this database are from North America, Europe, and developed countries in other regions, while small-

scale fisheries and those in developing countries, are underrepresented.

The second source of information used to infer fisheries stock status around the world comes from the United Nations Food and Agriculture Organization (FAO) landings database. The FAO conducts a regular stock status review (FAO 2014) focusing on biological overexploitation as defined in most fishery-related international treaties (e.g., United Nations Fish Stocks Agreement) and classifies stocks into three categories based on expert opinions: underexploited, fully exploited (including stocks that are notionally within 20% above or below the biomass that would support maximum sustainable yield, B_{MSY}), and overexploited. While the FAO stock status review includes more stocks than the RAM Legacy database, therefore providing a more comprehensive picture of stock status, the methods used to evaluate fisheries are more heterogeneous and less transparent than the stock assessments found in the RAM Legacy database. In particular, the reliance on expert judgment to determine these status estimates means that they are harder to reproduce, even with a systematic methodology (FAO 2014). Similarly, the estimates are categorical, which means they are less quantitative than those from a traditional stock assessment and may provide limited guidance for management decisions.

There are two major ways to improve estimates of individual stock status at the global level: (1) through improvements to the data and technical capacity available to quantitatively assess fisheries at the local level, and (2) through further refinements to methodologies that use widely available data (e.g., globally available landings data). The first option requires substantial new resources and expertise. The second option is more immediately attainable and a range of new approaches has recently been developed (e.g., Vasconcellos & Cochrane 2005; Berkson *et al.* 2011; Costello *et al.* 2012; Carruthers *et al.* 2014), which have fewer data requirements than traditional quantitative stock assessments. Here we focus on these approaches, which require a time series of removals (catches) and basic life history information. This enables us to obtain more complete global coverage of stock status than more data-intensive quantitative stock assessment models, although these methods are still subject to many limitations.

In particular, estimates of exploitation status by catch-only methods can be biased and highly uncertain largely due to required simplifying assumptions. Rosenberg *et al.* (2014) demonstrated the strengths and weaknesses of four catch-only models using a full factorial simulation framework, which evaluated their performance under different combinations of life-history traits, initial

depletion, effort dynamics, and length of the catch time series.

A recent development in data-limited stock assessment methods has been to combine the estimates of exploitation status, B/B_{MSY} , where B is the current stock biomass and B_{MSY} is the biomass estimated to result in maximum sustainable yield from multiple catch-only models in a superensemble (Anderson *et al.* In Press and Supporting Information). A superensemble (Krishnamurti *et al.* 1999) is “super” because it combines the estimates from multiple models by calibrating them on a data set with known or trusted properties—in this case the simulated data set from Rosenberg *et al.* (2014). Estimates from multiple models, here of B/B_{MSY} , are calibrated to known values via a regression model. This allows for both weighting the individual models based on their accuracy and exploiting the covariance between individual models to generate more accurate and less biased estimates of B/B_{MSY} . The superensemble approach is common in the weather and climate forecasting where, for example, it has been used to improve predictions of wind and precipitation in Asian monsoons (Krishnamurti *et al.* 1999) and improve global surface temperature forecasts (Berliner & Kim 2008).

Here, we quantitatively estimate current biomass of global fisheries relative to B_{MSY} for stocks in the FAO database. We aim to provide a more comprehensive assessment of global and regional stock status by implementing a superensemble approach to combine status estimates obtained from four different catch-only models.

Methods

We applied the four catch-only methods tested in Rosenberg *et al.* (2014) to 785 FAO stocks, subject to the criteria described in the Supporting Information (Table 1). We applied one empirical model, the panel regression approach (PRM) developed by Costello *et al.* (2012), and three mechanistic models, which all assume the same population dynamics, but make different assumptions about uncertainty and the dynamics of fishing effort:

- (1) the catch-MSY (C_{MSY}) model of Martell & Froese (2012), which includes assumptions about resilience;
- (2) the catch-only-model with sampling-importance resampling (COMSIR) developed by Vasconcellos & Cochrane (2005); and
- (3) the state-space catch-only model (SSCOM) developed by Thorson *et al.* (2013).

Further details on the models are presented in the Supporting Information. To obtain global estimates of stock status for all stocks, we combined the estimates from the four models using a superensemble (Supporting Information). The superensemble method goes well be-

yond simply averaging across individual method estimates. Our superensemble calibrated a combination of the four individual model estimates of B/B_{MSY} via a random forest method (Breiman 2001) fitted to a data set of nearly 6,000 simulated fish stocks with known B/B_{MSY} (Anderson *et al.* In Press). Random forests are a machine learning approach that allow for nonlinear relationships between the predictors (the individual model estimates) and the response (the superensemble estimate) and interactive effects between the individual model estimates while being relatively robust to overfitting (Breiman 2001). Previous analyses showed that a random forest superensemble consistently had the best or among the best performance characteristics when compared to other possible superensemble regression models (Anderson *et al.* In Press). The superensemble outperformed the individual models in cross-validation on simulated data with, for example, a median absolute proportional error in B/B_{MSY} of 0.32 compared to 0.42–0.56 for the individual models (Anderson *et al.* In Press).

Estimating stock status

We computed density plots to explore the distribution of stock status globally and within each FAO statistical region through 2013. We also compared our global estimates of status to other global estimates of status (see Table S2). Additionally, we compared our approach to traditional stock assessment estimates by matching stocks in the RAM Legacy database with those in the FAO catch database where possible. In some cases, there are multiple RAM stocks that match a single “stock” from the FAO database (e.g., tuna stocks or Atlantic cod). In these cases, we matched the RAM stock status estimate to each FAO region to which it could logically correspond. We also compared the status estimates for RAM Legacy-assessed stocks with the status of previously unassessed stocks from the FAO catch database.

Results

Global patterns

At the aggregate global level, the median B/B_{MSY} status of exploited stocks is 0.94, such that 439 stocks (56%) are estimated to be below the B_{MSY} reference point based on a superensemble of data-limited models (Table 1, Figure 1). Of these, 261 (59% of those below B_{MSY}) are estimated to be below 80% of the B_{MSY} level, which is the FAO (State of World Fisheries and Aquaculture) SOFIA definition of “overexploited.” Therefore, for the 178 stocks between B_{MSY} and 80% of B_{MSY} , significant yield may be foregone, but the current advice under FAO is that they are fully

Table 1 Numbers of “stocks” (i.e., fished taxa) per FAO statistical area under each data filter applied

FAO Area	Original # Stocks in FAO: 3,630 total	Filter: Species level taxa: 2,621 total	Filter: >20 years and >1,000 t of catch: 785 stocks	Median B/B_{MSY} for stocks in analysis	Percentage for stocks with $B/B_{MSY} < 1$ in analysis
Percentage of total FAO catches	100%	69.9%	66.1%	0.97	53%
Arctic Sea	7	5	NA	NA	NA
Atlantic, Antarctic	40	29	NA	NA	NA
Atlantic, Eastern Central	281	176	63	1.05	0.44
Atlantic, Northeast	379	269	96	0.95	0.57
Atlantic, Northwest	211	158	60	0.82	0.70
Atlantic, Southeast	194	122	33	0.84	0.76
Atlantic, Southwest	250	161	54	1.03	0.43
Atlantic, Western Central	247	157	52	0.84	0.67
Indian Ocean, Antarctic	51	34	NA	NA	NA
Indian Ocean, Eastern	219	108	55	1.13	0.29
Indian Ocean, Western	322	202	40	1.14	0.33
Mediterranean and Black Sea	255	163	51	1.02	0.47
Pacific, Antarctic	24	13	NA	NA	NA
Pacific, Eastern Central	190	111	31	1.00	0.52
Pacific, Northeast	110	80	26	0.98	0.54
Pacific, Northwest	223	131	90	0.82	0.67
Pacific, Southeast	203	121	46	0.91	0.59
Pacific, Southwest	209	126	38	1.02	0.47
Pacific, Western Central	215	96	50	1.10	0.34

Median B/B_{MSY} and probability of B/B_{MSY} pertain to stocks with species-level taxonomic resolution and with catch time series longer than 20 years and more than 1,000 t of annual median catch.

exploited, and unfortunately, no real policy change would be called for.

The superensemble was employed to deal with individual biases in each of the models, but underlying patterns could still be detected. For example, in addition to a primary mode below B/B_{MSY} , the superensemble estimated many stocks to have a B/B_{MSY} above 1, producing a second mode at 1.25. We investigated the distributions of the underlying data-limited models to understand the cause of this bimodality and found that the C_{MSY} and COMSIR models were mainly responsible for this pattern in the estimates. In a sensitivity analysis for C_{MSY} , we found that the bimodality was due to the prior distributions assigned to the final year depletion, which are based on the catch trajectories (Figure S3). This bimodality carries forward into the superensemble estimates. However, the overall results do not change if each model is removed individually from the superensemble (see Supporting Information Section S1.5, *Sensitivity analyses*). The SSCOM method frequently estimated stocks to be underexploited relative to B_{MSY} (Figure 1). The ensemble partly accounts for potential systematic bias through the relative weightings, but these estimates of higher biomass still affect the overall pattern.

Regional patterns

For 9 of the 15 FAO regions, a majority of the stocks were estimated to be below B_{MSY} (Table 1, Figure 2). This is particularly striking in the Northern Hemisphere regions, where most stocks are estimated to be below the biomass that would support MSY (Figure 2). The exception to this pattern was the NE Pacific, where more stocks were above B_{MSY} . Similarly, the majority of stocks in the South Atlantic and South Pacific regions were below B_{MSY} .

In the Atlantic, all of the areas have a modal value for estimated stock status below 80% of B_{MSY} (Figure 2) but with substantial variation in status among stocks within each region. In all of the Atlantic regions with the exception of the Eastern Central, the majority of the stocks are below B_{MSY} (Figure 2). Several Atlantic regions show a bimodal distribution of stock status.

Similarly, the NW Pacific, SW Pacific, and SE Pacific regions have the majority of stocks below B_{MSY} while the NE Pacific and Western Central Pacific have slightly more stocks above B_{MSY} than below (Figure 2). The Indian Ocean's two regions are in better condition with only about one-third of the stocks below B_{MSY} . The long tails on the distribution of stock status for all regions




Figure 1 Global distribution of B/B_{MSY} status for 785 analyzed stocks estimated by each data-limited method and the superensemble approach.

indicates that there are some stocks that are only lightly exploited, at least with regard to recent catch trends.

Critically, within all of the regions a substantial number of stocks are estimated to be within 20% below B_{MSY} such that they would be classed as “fully exploited” in previous studies.

Comparisons with other methods

Direct comparison of the ensemble estimates for stocks that are also included in the RAM Legacy database (i.e., a comparison of assessed RAM to assessed ensemble stocks) show that for most regions our methods are slightly more pessimistic concerning stock status than analytical stock assessments (Figure S4). This is also the case when comparing estimates of the status of assessed stocks contained in the RAM Legacy database with estimates using the ensemble method for those stocks without analytical

assessments contained in the FAO data we utilized (Figure 3; i.e., a comparison of assessed RAM to unassessed stocks). However, many more stocks can be considered using the catch-only methods than can be analytically assessed by traditional methods.

Discussion

Our results suggest that the overall status of fisheries globally is below the biomass capable of producing maximum sustainable yield (Table 1). However, according to the FAO SOFIA definitions and methodology of expert opinion, 41% of these stocks would have been classified as “fully exploited,” rather than “overexploited.” From a manager’s point of view, perhaps no action would be needed to improve management of a fully exploited stock, under the FAO SOFIA categorical status assessments.




Figure 2 Regional distribution of B/B_{MSY} status for analyzed stocks estimated by the superensemble approach. Black vertical lines indicate $B/B_{MSY} = 1$ with dark gray bars indicating $\pm 20\%$ around this point. The density curves have been truncated at $B/B_{MSY} = 2$ for ease of interpretation. This truncation results in at least 98.75% of the estimates captured by the area under the curve. The panels are organized in order of decreasing median B/B_{MSY} of the superensemble.

However, the more quantitative estimates provided in this study highlight the possible need for management action for many of these taxa.

Our results overall are broadly consistent with other studies that have attempted to provide a global picture of fishery status using a variety of methods (Table S2). A key advantage of using this approach is that it does not require a wholesale change in estimation method each time a new method becomes available. Our methodology can be easily repeated as new information becomes available, it is objective, and can incorporate new methods as part of the superensemble. Combining estimates from dif-

ferent methods in a consistent reproducible manner may provide more stability in the advice for managers. Additionally, the superensemble approach is more robust than typically model averaging because the superensemble does not simply average across individual method estimates of status but uses those estimates as input to a new statistical model that is then trained on known status, which is a key advantage of this approach.

Recent requirements to set scientifically based catch limits in several countries (Rosenberg *et al.* 2009) and growing consumer demand for sustainably managed seafood products (Gutierrez *et al.* 2012) have spurred




Figure 3 Comparison of B/B_{MSY} estimates by FAO area for unassessed versus assessed stocks. Assessed B/B_{MSY} was calculated as the median of the B/B_{MSY} estimates from stocks in the RAM Legacy database. Unassessed B/B_{MSY} was calculated as the median of the B/B_{MSY} estimates determined by the superensemble approach for all stocks without a RAM database analog. The estimates are ordered by decreasing median assessed B/B_{MSY} estimates.

an emerging field of methods for estimating overfishing thresholds and setting catch limits for stocks with limited data (e.g., Berkson *et al.* 2011; Carruthers *et al.* 2014). Stock status is not the sole input guiding management, but it should provide a key indication of whether a stock is in a safe operating space. Indeed, there is broad consensus among management authorities worldwide that managing stocks toward values within the range of MSY is an important guidepost for achieving fisheries sustainability as shown by the UN Fish Stocks Agreement under the Law of the Sea.

Often, the status of quantitatively assessed stocks is used to infer the status of unassessed stocks. However,

several studies (Worm *et al.* 2009; Costello *et al.* 2012; Ricard *et al.* 2012; Costello *et al.* 2016) have shown that this inference may be seriously flawed. In consequence, continued overexploitation of unassessed stocks, often by small-scale fisheries, has led to significant loss in benefits to people (Inniss *et al.* 2016). Our results are generally more pessimistic regarding stock status than full stock assessments (Figure 3 and Figure S4). These differences may be due to better management of assessed stocks as well as any methodological differences, though ascribing the effect to either cause is not a simple matter. We recognize that our methods, and all stock assessments, are limited by the time series of catch information

available and the dynamics of that catch. If large declines in abundance occurred prior to the start of the data series for a given stock an incorrect impression of stock status will obtain.

Although our approach provides an important step toward quantitative and replicable estimates of stock status for a larger set of fished stocks than has been the case in the past, there are still many limitations to using this information for stock-specific or even regional advice. These include the high variability of the estimates, the need for longer time series of data, limited life history information for many stocks, and the difficulties of assigning prior distributions, particularly given the problem of bimodality in some of the distributions of our estimates. The simulation testing described in the Supporting Information and Rosenberg *et al.* (2014) can help explore these issues but does not resolve them. There are fundamental data limitations that are inherent to the problem we are addressing. Key potential areas for future improvement include better informed prior distributions for final status in different regions, including priors that account for the characteristics of fishery resources in different regions (Cope *et al.* 2015), inclusion of existing survey and fishery data (Thorson *et al.* 2012), and evaluation of how well these models inform management strategies for data-limited fisheries.

Acknowledgment

This research was funded by the Food and Agricultural Organization of the United Nations, Conservation International and the Gordon and Betty Moore Foundation.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web site:

Table S1. Summary of variables used in the modified panel regression model (mPRM)

Table S2. Comparison of estimates of stock status from various recent studies illustrating the different definitions and numbers of "stocks" evaluated

Figure S1. Individual model and spectral covariate contributions to the random forest superensemble (random forest partial dependence plots).

Figure S2. Violin plots showing median, interquartile range, and distribution of B/B_{MSY} estimates from ensemble models that exclude each model in turn, i.e., each data-limited model and the spectral densities of the catch trajectories, showing how the removal of each covariate influences the overall global status estimates, compared to the baseline model with all covariates retained: "all."

Figure S3. Illustration of the variability in the B/B_{MSY} estimates from C_{MSY} when the priors on initial and final depletion are varied, with an example of stocks from the Pacific Southwest and Pacific Northeast FAO regions.

Figure S4. Comparison of B/B_{MSY} estimates by FAO area for assessed RAM stocks and assessed stocks in the ensemble.

References

- Anderson, S.C., Branch, T.A., Ricard, D. & Lotze, H.K. (2012). Assessing global marine fishery status with a revised dynamic catch-based method and stock-assessment reference points. *ICES J. Marine Sci.*, **69**, 1491-1500.
- Anderson, S.C., Cooper, A.B., Jensen, O.P., *et al.* (In Press). Improving estimates of population status and trend with superensemble models. *Fish and Fisheries*.
- Berkson, J., Barbieri, L., Cadrin, S. *et al.* (2011). Calculating acceptable biological catch for stocks that have reliable catch data only (Only Reliable Catch Stocks - ORCS). p. 56. *NOAA Technical Memorandum NMFS-SEFSC-616*. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southeast Fisheries Science Center, Miami, FL.
- Berliner, L.M. & Kim, Y. (2008). Bayesian design and analysis for superensemble-based climate forecasting. *J. Climate*, **21**, 1891-1910.
- Beverton, R.J.H. & Holt, S.J. (1957). *On the dynamics of exploited fish populations*. Fishery Investigations, London.
- Breiman, L. (2001). Random forests. *Machine Learning*, **45**, 5-32.
- Carruthers, T., Punt, A.E., Walters, C.J. *et al.* (2014). Evaluating methods for setting catch limits in data-limited fisheries. *Fisheries Res.*, **153**, 48-68.
- Cope, J.M., Thorson, J.T., Wetzell, C.R. & DeVore, J. (2015). Evaluating a prior on relative stock status using simplified age-structured models. *Fisheries Res.*, **171**, 101-109.
- Costello, C., Ovando, D., Clavelle, T. *et al.* (2016). Global fishery prospects under contrasting management regimes. *Proc. Natl. Acad. Sci.*, **113**, 5125-5129.
- Costello, C., Ovando, D., Hilborn, R., Gaines, S.D., Deschenes, O. & Lester, S.E. (2012). Status and solutions for the world's unassessed fisheries. *Science*, **338**, 517-520.
- FAO. (2014). The State of World Fisheries and Aquaculture 2014. Page 223 in FAO, editor. Food and Agriculture Organization of the United Nations, Rome.
- Fernandes, P.G. & Cook, R.M. (2013). Reversal of fish stock decline in the Northeast Atlantic. *Curr. Biol.*, **23**, 1432-1437.
- Garcia, S.M. & Rosenberg, A.A. (2010). Food security and marine capture fisheries: characteristics, trends, drivers and future perspectives. *Philos. T. Roy. Soc. B*, **365**, 2869-2880.
- Gutierrez, N.L., Valencia, S.R., Branch, T.A. *et al.* (2012). Eco-label conveys reliable information on fish stock health to seafood consumers. *Plos One*, **7**.

- Hilborn, R. & Ovando, D. (2014). Reflections on the success of traditional fisheries management. *ICES J. Marine Sci.*, **71**, 1040-1046.
- Hilborn, R. & Walters, C. (1992). *Quantitative fisheries stock assessment: choice, dynamics and uncertainty*. Chapman and Hall, New York.
- Inniss, L., Simcock, A., Ajawin, A.Y. *et al.* (2016). *The first global integrated marine assessment; world ocean assessment*. United Nations, New York.
- Krishnamurti, T.N., Kishtawal, C.M., LaRow, T.E. *et al.* (1999). Improved weather and seasonal climate forecasts from multimodel superensemble. *Science*, **285**, 1548-1550.
- Martell, S.J. & Froese, R. (2012). A simple method for estimating MSY from catch and resilience. *Fish Fisheries*, **14**, 504-514.
- Pauly, D., Christensen, V., Dalsgaard, J., Froese, R. & Torres, Jr. F. (1998). Fishing down marine food webs. *Science*, **279**, 860-863.
- Quinn, T.J. & Deriso, R.B. (1999). *Quantitative fish dynamics*. Oxford University Press, New York.
- Ricard, D., Minto, C., Jensen, O.P. & Baum, J.K. (2012). Examining the knowledge base and status of commercially exploited marine species with the RAM Legacy Stock Assessment Database. *Fish Fisheries*, **13**, 380-398.
- Rosenberg, A.A., Acosta, A., Babcock, E. *et al.* (2009). Use of productivity-susceptibility analysis (PSA) in setting annual catch limits for U.S. fisheries: a workshop report. Page 18 in I. MRAG Americas, editor. Lenfest Ocean Program, Washington, DC.
- Rosenberg, A.A., Fogarty, M.J., Cooper, A.B. *et al.* (2014). Developing new approaches to global stock status assessment and fishery production potential of the seas. p. 175. *FAO Fisheries and Aquaculture Circular No 1086*. FAO, Rome.
- Thorson, J.T., Branch, T.A. & Jensen, O.P. (2012). Using model-based inference to evaluate global fisheries status from landings, location, and life history data. *Canadian J. Fisheries Aquat. Sci.*, **69**, 645-655.
- Thorson, J.T., Minto, C., Minto-Vera, C.V., Kleisner, K. & Longo, C. (2013). A new role for effort dynamics in the theory of harvested populations and data-poor stock assessment. *Canadian J. Fisheries Aquat. Sci.*, **70**, 1829-1844.
- Vasconcellos, M. & Cochrane, K. (2005). Overview of world status of data-limited fisheries: inferences from landings statistics. Pages 1-20 in G.H. Kruse, V.F. Gallucci *et al.*, editors. *Fisheries assessment and management in data-limited situations; Proceedings: Lowell Wakefield Symposium, 21, Anchorage, AK (USA), 22-25 Oct 2003*. Alaska Sea Grant College Program, Fairbanks.
- Worm, B., Hilborn, R., Baum, J.K. *et al.* (2009). Rebuilding global fisheries. *Science*, **325**, 578-585.